

From Wigner–Inönü Group Contraction to Contractions of Algebraic Structures

N.A. Gromov

Department of Mathematics, Institute of Mathematics and Mechanics,
Ural Division, Russian Academy of Sciences, Syktyvkar 167982, Russia

E-mail: gromov@dm.komisc.ru

Abstract

The development of the notion of group contraction first introduced by E. Inönü and E.P. Wigner in 1953 is briefly reviewed. The fundamental role of the idea of degenerate transformations is stressed. The significance of contractions of algebraic structures for exactly solvable problems of mathematical physics is noticed.

1 Introduction

Starting with the fact that classical mechanics is a limiting case of relativistic mechanics and therefore the Galilei group must be in some sense a limiting case of the relativistic mechanics' group E.P. Wigner and E. Inönü in 1953 have introduced [1] a new notion in group theory, namely, a *contraction* of group (Lie algebra and their representations) as follows: "We shall call the operation of obtaining a new group by a singular transformation of the infinitesimal elements of the old group a *contraction* of the latter." This concept is based on the principal idea: only singular transformation of Lie algebra gives in result a new nonisomorphic algebra.

The concept of contraction turned out to be fruitful and was formulated [2] as the contraction of an arbitrary algebraic structure $(M, *)$ with the help of ϵ -depending map

$$\phi_\epsilon : (M, *) \rightarrow (N, *'),$$

where $(N, *')$ is the algebraic structure of the same type, isomorphic to $(M, *)$ for $\epsilon \neq 0$ and nonisomorphic to the initial one for $\epsilon = 0$. Except for original Wigner–Inönü group (algebra) contractions the contractions of Lie superalgebra was introduced [3] and after that more general graded contractions [4], [5] which preserve an arbitrary grading of Lie algebra and may be applied to an infinite-dimensional algebras [6]. Under contractions of bialgebra [7] Lie algebra structure and cocommutator are conserved. Hopf algebra (quantum algebra [8], [9] and quantum group [10]) contractions are introduced in such a way, that in

the limit $\epsilon \rightarrow 0$ a new expressions for coproduct, counit and antipode are consistent with the Hopf algebra axioms. Recently contractions of the algebraic structures with bilinear products of arbitrary nature on sections of finite-dimensional vector bundles was presented [11].

2 Contractions of algebraic structures

A **Lie algebra** L is a vector space V equipped with a Lie bracket $[\cdot, \cdot] : L \otimes L \rightarrow L$, which is linear in both arguments, antisymmetric and satisfy the Jacobi identity. So the structure of Lie algebra is as follows: $L = (V, [\cdot, \cdot])$. It is well known fact that the non-degenerate linear transformations $\Phi : V \rightarrow V$, $\det \Phi \neq 0$ of vector space V give in result the Lie algebra L' which is isomorphic to the initial one's L . How to receive a new Lie algebra from L ? The answer was given fifty years ago by E. İnönü and E.P. Wigner [1]: it is necessary to use a degenerate transformation of V . Let us regard $\Phi_\epsilon : V \rightarrow V$ such that $\det \Phi_\epsilon \neq 0$ for $\epsilon \neq 0$, but $\det \Phi_\epsilon = 0$ for $\epsilon = 0$. Then if the limit

$$[\cdot, \cdot]' = \lim_{\epsilon \rightarrow 0} \Phi_\epsilon \circ [\cdot, \cdot] \circ (\Phi_\epsilon^{-1} \otimes \Phi_\epsilon^{-1}) \quad (1)$$

exist, we have the new Lie algebra $L' = (V, [\cdot, \cdot]')$ with the same underlying vector space. The limit (1) is called a *contraction of Lie algebra*.

A **Lie superalgebra** has the structure $L_s = (V = V_0 \oplus V_1, < \cdot, \cdot >, Z_2) = L_s^0 \oplus L_s^1$, where $< \cdot, \cdot >$ is a bilinear product (commutator or anticommutator), Z_2 is a grading group, such that V_0 and V_1 are even and odd parts of vector space V , respectively, L_s^0 and L_s^1 are even and odd parts of superalgebra L_s . The nondegenerate limit [3]

$$< \cdot, \cdot >' = \lim_{\epsilon \rightarrow 0} \Phi_\epsilon \circ < \cdot, \cdot > \circ (\Phi_\epsilon^{-1} \otimes \Phi_\epsilon^{-1}), \quad (2)$$

where Φ_ϵ preserves Z_2 -grading, gives in result contracted superalgebra $L'_s = (V = V_0 \oplus V_1, < \cdot, \cdot >', Z_2) = L_s^{0'} \oplus L_s^{1'}$.

Graded contractions were introduced in [4], [5] and deal with the structure (L, Γ) , where L is a Lie algebra and Γ is a finite Abelian group. A grading of L by Γ means the decomposition of the vector space V of L into a direct sum of subspaces

$$V = \bigoplus_{\mu \in \Gamma} V_\mu \quad (3)$$

such that for every choice of $X \in V_\mu$, $Y \in V_\nu$, if $[X, Y] \neq 0$, then $[X, Y] \in V_{\mu+\nu}$. Symbolically

$$0 \neq [V_\mu, V_\nu] \subseteq V_{\mu+\nu}, \quad \mu, \nu, \mu + \nu \in \Gamma. \quad (4)$$

In this case the map Φ_ϵ additionally to existence of (1) has to preserve the grading equations (3), (4). Graded contractions leads to a structure $(L', \Gamma) = (V, [\ , \], \Gamma)$ with the same underlying V and the same grading group Γ . Graded contractions of the inifitedimensional Virasoro algebras were regarded in [6].

A **Lie bialgebra** (L, η) is a Lie algebra L endowed with a cocomutator $\eta : L \rightarrow L \otimes L$ such that

i) η is 1-cocycle, i.e.

$$\eta([X, Y]) = [\eta(X), 1 \otimes Y + Y \otimes 1] + [1 \otimes X + X \otimes 1, \eta(Y)], \quad \forall X, Y \in L, \quad (5)$$

ii) the dual map $\eta^* : L^* \otimes L^* \rightarrow L^*$ is a Lie bracket on L^* .

Contracted Lie brackets are obtained by (1). Contraction of cocommutator η may be realized by the mapping $\Psi_\epsilon : L \rightarrow L'$, which is in general different from Φ_ϵ , as follows:

$$\eta' = \lim_{\epsilon \rightarrow 0} (\Psi_\epsilon \otimes \Psi_\epsilon) \circ \eta \circ \Psi - \epsilon^{-1}. \quad (6)$$

The consistency of $[\ , \]$ and η gives the another expression for cocommutator

$$\eta' = \lim_{\epsilon \rightarrow 0} (\Psi_\epsilon \otimes \Psi_\epsilon) \circ \eta \circ \Phi_\epsilon^{-1}. \quad (7)$$

If all limits (1), (6), (7) exist, then (L', η') is a *contracted Lie bialgebra* [7].

An associative algebra A is said to be a **Hopf algebra** if there exist maps: coproduct $\Delta : A \rightarrow A \otimes A$, counit $u : A \rightarrow \mathbb{C}$ and antipode $\gamma : A \rightarrow A$, such that $\forall a \in A$ the following axiomas are hold

$$\begin{aligned} H_1 : \quad & (id \otimes \Delta)\Delta(a) = (\Delta \otimes id)\Delta(a), \\ H_2 : \quad & (id \otimes u)\Delta(a) = (u \otimes id)\Delta(a), \\ H_3 : \quad & m((id \otimes \gamma)\Delta(a)) = m((\gamma \otimes id)\Delta(a)), \end{aligned} \quad (8)$$

where $m(a \otimes b) = ab$ is the usual multiplication. So the Hopf algebra structure is $H = (A, \Delta, u, \epsilon)$. Contractions of coproduct, counit and antipode are realized by the same mapping χ_ϵ of A . Therefore, if the limits

$$\Delta' = \lim_{\epsilon \rightarrow 0} (\chi_\epsilon \otimes \chi_\epsilon) \circ \Delta \circ \chi_\epsilon^{-1}, \quad u' = \lim_{\epsilon \rightarrow 0} u \circ \chi_\epsilon^{-1}, \quad \gamma' = \lim_{\epsilon \rightarrow 0} \chi_\epsilon \circ \gamma \circ \chi_\epsilon^{-1}. \quad (9)$$

exist, then $H' = (A, \Delta', u', \epsilon')$ is a *contracted Hopf algebra*. Contractions of Hopf algebras on the level of quantum algebras were first developed in [8],[9] and on the level of quantum groups in [10].

The general conception of contraction of algebraic structures was recently developed [11] to contractions of $(A, *)$, where A is a set of

smooth sections of vector bundle E over a manifold M , $*$: $A \times A \rightarrow A$ is a point-wise continuous bilinear product are given by the following

Theorem. Let $N : E \rightarrow E$ be a smooth $(1,1)$ -tensor. Denote by $U(\epsilon) = \epsilon I + N$ a deformation of N , by $E = E^1 \oplus E^2$ the Riesz decomposition of E relative to N , and by A^1 the set of smooth sections of E^1 . Then the limit

$$X *_N Y = \lim_{\epsilon \rightarrow 0} U^{-1}(\epsilon)(U(\epsilon)(X) * U(\epsilon)(Y)) \quad (10)$$

exists for all $X, Y \in A$ and defines a new (contracted) bilinear product on A if and only if the torsion

$$T_N(X*Y) = N(X)*N(Y) - N(N(X)*Y + X*N(Y)) + N^2(X*Y) \quad (11)$$

takes values in $N(A^1)$.

Lie algebras, Lie algebroids and Poisson brackets were given as an examples of the developed theory. In particular, Wigner-Inönü Lie algebra contractions correspond to the vector bundle E over a single point with $A = E$, $*$ = $[,]$, $U(\epsilon) = P + \epsilon(I - P)$, where P is a projection.

3 Conclusions

The notion of contraction first introduced by E. Inönü and E.P. Wigner for Lie groups and algebras is now extend on new algebraic structures. Some of them are mention in this talk. But the fundamental idea of degenerate transformation is presented in all cases.

It is well known that often a symmetry is a necessary condition for explicit solutions of many mathematical and physical problems. And at the second half of the last century a models of the mathematical physics are usually formulated with the help of different mathematical structures (Lie groups and algebras, superalgebras, quantum groups etc.), which are a mathematical tools for describing symmetries. Above-mentioned algebraic structures constitute the base of exactly solvable problems of mathematical physics and its contractions are correspond to different limiting cases of these problems.

References

- [1] E. Inönü and E.P. Wigner, *Proc. Nat. Acad. Sci. USA* **39** (1953) 510.
- [2] N.A. Gromov, *Turkish J. Phys.*, **21** (1997) 377.

- [3] J. Rembielinski and W. Tybor, *Acta Phys. Polonica* **B15** N 7 (1984) 611.
- [4] M. de Montigny and J. Patera, *J. Phys. A: Math. Gen.* **24** (1991) 525.
- [5] R.V. Moody and J. Patera, *J. Phys. A: Math. Gen.* **24** (1991) 2227.
- [6] I.V. Kostyakov, N.A. Gromov and V.V. Kuratov, *Nucl. Phys. B (Proc. Suppl.)*, **102&103** (2001) 316.
- [7] A. Ballesteros, N.A. Gromov, F.J. Herranz, M.A. del Olmo and M. Santander, *J. Math. Phys.* **36** (1995) 5916; hep-th/9412083.
- [8] E. Celeghini, R. Giachetti, E. Sorace and M. Tarlini, *J. Math. Phys.* **32** (1991) 1155.
- [9] E. Celeghini, R. Giachetti, E. Sorace and M. Tarlini, *J. Math. Phys.* **32** (1991) 1159.
- [10] N.A. Gromov, *J. Phys. A: Math. Gen.* **26** (1993) L5.
- [11] J.F. Cariñena, J. Grabowski and G. Marmo, *J. Phys. A: Math. Gen.* **34** (2001) 3769.